

Journal of Physics B: Atomic, Molecular and Optical Physics



PAPER

OPEN ACCESS

RECEIVED
5 February 2026

REVISED
23 April 2026

ACCEPTED FOR PUBLICATION
5 June 2026

PUBLISHED
2 July 2026

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Retrieval of the nuclear motion in a molecule from photoelectron momentum distributions using non-Born–Oppenheimer quantum dynamics and deep learning

N I Shvetsov-Shilovski^{1,2,*} and M Lein²

¹ Rudolf Bembeneck Gesamtschule Burgdorf, Vor dem Celler Tor 50, 31303 Burgdorf, Germany

² Institut für Theoretische Physik, Leibniz Universität Hannover, 30167 Hannover, Germany

* Author to whom any correspondence should be addressed.

E-mail: n79@narod.ru

Keywords: time-dependent bond-length, deep learning, strong-field ionization, photoelectron momentum distributions

Abstract

By using a neural network that takes momentum distributions of photoelectrons produced in strong-field ionization as input, we retrieve the time-dependent bond length of a dissociating one-dimensional H_2^+ molecule. The photoelectron momentum distributions are calculated from the direct numerical solution of the non-Born–Oppenheimer time-dependent Schrödinger equation. We simulate two setups: first, molecules prepared in the first excited electronic state, second, a pump-probe scheme starting from the ground state. We show that in both schemes a neural network trained on momentum distributions calculated for frozen nuclei retrieves the time-dependent bond length with an absolute error of $0.2 - 0.4$ a.u. We find that a neural network, when applied to photoelectron momentum distributions obtained within the pump-probe scheme, can be used for the retrieval of the equilibrium internuclear distance and ground-state population. This opens new perspectives for extracting electronic properties of molecules from electron momentum distributions using deep learning.

1. Introduction

The past few years have witnessed remarkable progress in the development of artificial intelligence and its subfield machine learning (ML). ML is currently widely used in all branches of physics: from astrophysics to soft-matter, medical, and biological physics, see, e.g., [1, 2] for reviews. This is due to the fact that ML is capable of processing huge amounts of data and drawing conclusions based on these data. Furthermore, ML methods are used in order to find complex mappings that depend on many variables and are hidden in the input data.

ML has also begun to be used in strong-field physics, which studies the interaction of strong laser radiation with atoms and molecules. The interaction of quantum systems with intense laser pulses results in a variety of highly non-linear phenomena, including above-threshold ionization (ATI), high-order harmonic generation (HHG), and sequential and nonsequential double ionization, see, e.g., [3–8] for reviews. Among applications of ML in strong-field physics are the prediction of the flux of high-order harmonics [9], reconstruction of the intensity and the carrier-envelope phase of short laser pulses from 2D images (frequency-resolved optical gating traces [10] and dispersion scan traces [11]), the efficient implementation of the trajectory-based Coulomb-corrected strong-field approximation (TCSFA) [12] (the TCSFA method is developed in [13, 14]), the retrieval of the femtosecond pulse parameters based on photoelectron momentum distributions (PMDs) produced in the ATI process [15], and the simulation of the macroscopic HHG driven by structured laser beams [16]. Another remarkable example is the prediction of HHG spectra for model diatomic and triatomic molecules for randomly chosen laser intensity, internuclear distance, and orientation of the molecule using ML, as well as the solution of the inverse problem: retrieval of the molecular and laser parameters and determination of the type of molecule based on the high-harmonic spectrum or time-dependent dipole acceleration [17]. Recently,

ML methods have been applied to the identification of the geometrical structure of molecules in laser-induced electron diffraction (LIED) [18], creating movies of attosecond charge migration using high harmonic spectroscopy [19], and the determination of the internuclear distance in a molecule from a given PMD [20–22]. These latter applications can be used as tools for time-resolved molecular imaging.

Time-resolved molecular imaging aims to visualize electronic and molecular dynamics in real time. The development of such techniques is extremely important for many branches of science and technology, including chemistry, biology and material science, as they provide a better understanding of various processes in molecules. Among the well-known methods of time-resolved molecular imaging are optical pump-probe spectroscopy, time-resolved x-ray absorption spectroscopy, time-resolved electron diffraction and time resolved x-ray diffraction, time-resolved scanning tunneling spectroscopy and others (see [23] for a review). Some of the processes studied in strong-field physics are applicable to molecular imaging. For example, the LIED (see, e.g., [24–26]) and strong-field photoelectron holography (SFPH) methods [27] analyze electron momentum distributions produced in the ATI process.

These methods are based on the fact that PMDs encode a lot of time-resolved information about the ion and the photoelectron. Two different kinds of electron wave packets generated in strong-field ionization contribute to the PMDs. The low-energy part of the distributions is formed to a large extent by direct electrons that do not return to their parent ions. In contrast to this, the high-energy part of PMDs is created by rescattered electrons that are driven back by the oscillating laser field to their parent ions and scatter from them by angles close to 180° . The LIED method retrieves static and time-resolved information from the high-energy part of PMD, i.e., from the rescattered electrons. In contrast to this, SFPH analyzes the holographic patterns emerging due to the interference of direct and rescattered electrons in the low-energy part of PMDs.

Both the LIED and SFPH techniques have been intensively developed over the last decade, see [28] for a recent review of the LIED method and [29–35] for the modern achievements in studies of SFPH. In [20–22] a convolutional neural network (CNN) was used for the retrieval of the static and time-dependent internuclear distance in a two-dimensional (2D) H_2^+ molecule based on the whole PMD (low and high-energy parts). It was shown that the CNN can retrieve the fixed internuclear distance with an absolute error of less than 0.1 a.u. [20]. In contrast to the method of [18], where a CNN was trained on the 2D differential cross sections, the information encoded in the electron momentum distributions was exploited directly in [20]. The corresponding PMDs were obtained from the numerical solution of the time-dependent Schrödinger equation (TDSE), and therefore a number of approximations that are usually made in the LIED and SFPH techniques were avoided. By using the transfer learning technique [36], the CNN [20] has been made applicable to distributions calculated for parameters it was not explicitly trained for, including focal-volume averaged PMDs and PMDs from another molecular species [21].

In [22] a CNN has been applied for reconstruction of the time-dependent bond length in the dissociating 2D H_2^+ molecule based on electron momentum distributions obtained within a pump-probe scheme. As in [20, 21], the PMDs were obtained from the solution of the TDSE, and the motion of the nuclei was treated classically, semiclassically and quantum mechanically. In all these cases, the neural network trained on PMDs calculated for frozen nuclei predicts the time-dependent internuclear distance with an absolute error of 0.2–0.3 a.u. However, the quantum approach of [22] is approximate: it does not account for the motion of the nuclei during the probe laser pulse. It is therefore of interest to investigate whether ML is able to reconstruct the time-dependent bond length from PMDs obtained by treating the dynamics of the molecule fully quantum mechanically without any approximations. In this paper we address the formulated problem by employing non-Born–Oppenheimer simulations. For simplicity, we consider a dissociating one-dimensional (1D) H_2^+ molecule. The 1D model restricts the motion of electrons and nuclei to a single spatial dimension, which leads to a number of limitations. First, the 1D model does not realistically describe the three-dimensional (3D) molecular orbitals, which can have angular momentum. Second, it does not accurately describe the 3D nature of molecular orbital overlap. This leads to inaccuracies in calculating bond lengths, dissociation energies, and vibrational frequencies. Nevertheless, the 1D model is useful as a qualitative model for understanding the basic physics of the phenomena under study.

We use two different schemes. First, we assume an instantaneous excitation of the molecule, i.e. the initial state is a nuclear wave packet prepared in the first excited electronic state. Second, we consider a pump-probe scheme starting from the molecular ground state, i.e., the molecule is excited by a pump pulse and ionized by a time-delayed probe pulse. We demonstrate that neural networks trained on PMDs calculated with moving nuclei in both schemes retrieve the internuclear distance with good accuracy. We then study the application of a neural network trained on momentum distributions calculated for frozen nuclei to the retrieval of the time-dependent internuclear distance. We find that in both instantaneous excitation and pump–probe schemes, a neural network trained on PMDs for fixed

nuclei can retrieve the time-dependent internuclear distance with an absolute error of about 0.2–0.4 a.u. Finally, we demonstrate that deep learning can be used for the reconstruction of not only nuclear motion but also the electronic properties of the molecule. In the pump–probe scheme, we use a neural network trained on PMDs for fixed nuclei to reconstruct the electronic ground-state population at the end of the pump pulse.

The paper is organized as follows. In section 2 we discuss the method used to solve the TDSE and to calculate the PMDs as well as the architecture of neural networks used in the present study. In section 3 we train and apply our neural networks to both schemes, and we analyze the obtained results. The conclusions are given in section 4. Atomic units are used throughout the paper unless indicated otherwise.

2. Methods

2.1. Solution of the TDSE

The time-dependent Schrödinger equation for a model H_2^+ molecule, in which the electronic and nuclear motions are restricted to one spatial dimension, is given by

$$i\frac{\partial}{\partial t}\Psi(x, R, t) = \left\{ -\frac{1}{2\mu}\frac{\partial^2}{\partial x^2} - \frac{1}{m_p}\frac{\partial^2}{\partial R^2} - i\frac{m_p+1}{m_p}A(t)\frac{\partial}{\partial x} + V_{pe}(x, R) + \frac{1}{\sqrt{R^2 + \alpha_1}} \right\} \Psi(x, R, t), \quad (1)$$

where x is the electron coordinate, R is the internuclear distance, $\Psi(x, R, t)$ is the wave function, m_p is the proton mass, $A(t)$ is the vector potential of the laser pulse, μ is the reduced mass for the electron motion, α_1 is the soft-core parameter for the proton–proton interaction, and the potential V_{pe} describes the interaction between the electron and both protons. It is worth noting that for a homonuclear diatomic molecule, the laser field does not couple to the R coordinate, see [37]. The velocity gauge is used in equation (1). The potential V_{pe} is chosen as

$$V_{pe}(x, R) = -\frac{1}{\sqrt{(x - \frac{1}{2}R)^2 + \alpha_2}} - \frac{1}{\sqrt{(x + \frac{1}{2}R)^2 + \alpha_2}}, \quad (2)$$

where α_2 is the soft-core parameter. Following [38], we have chosen $\alpha_1 = 0.03$ and $\alpha_2 = 1.0$.

The vector potential $A(t)$ of the ionizing laser pulse is given by

$$A_{\text{probe}}(t) = (-1)\frac{F_0}{\omega}f(t)\cos[\omega(t - t_c)], \quad (3)$$

where ω is the laser frequency, F_0 is the field-strength amplitude, $t = t_c$ is the center of the laser pulse, τ_L is the pulse duration, and the envelope $f(t)$ is given by

$$f(t) = \begin{cases} \sin^2 \left[\frac{\omega(t - t_c + \frac{\tau_L}{2})}{2n_p} \right] & \text{for } t_c - \frac{\tau_L}{2} \leq t \leq t_c + \frac{\tau_L}{2} \\ 0 & \text{otherwise.} \end{cases} \quad (4)$$

Here, n_p is the number of the optical cycles within the pulse and, therefore, $\tau_L = (2\pi/\omega) \cdot n_p$. We perform our simulations for $\varphi = 0$.

Equation (1) or similar equations has been used in many studies of H_2^+ interacting with strong laser pulses, see, e.g., [38–40]. In order to solve the TDSE (1), we use the Feit–Fleck–Steiger split-operator method [41]. The computational box extends over $x \in [-400, 400]$ a.u. and $R \in [0, 80]$ a.u. The grids in x and R directions consist of 4096 and 2048 points, respectively, which corresponds to the grid spacings $\Delta x = 0.1954$ a.u. and $\Delta R = 0.0391$ a.u. The TDSE is propagated in time with the time step $\Delta t = 0.0184$ a.u. We use absorbing boundaries in the x direction, in order to prevent unphysical reflections of the wave function from the boundary of the computational grid, i.e., at every time step we multiply the wave function by the mask:

$$M(x) = \begin{cases} 1 & \text{for } |x| \leq x_b \\ \exp[-\beta(x - x_b)^2] & \text{for } |x| > x_b, \end{cases} \quad (5)$$

where $\beta = 10^{-4}$ and $x_b = 300$ a.u. The potential V_{pe} is neglected for $|x| > x_b$, but the interaction between the protons described by the term $1/\sqrt{R^2 + \alpha_1}$ term in equation (1), is taken into account in

this region. Indeed, large values of x do not necessarily imply large values of R . Due to the large extent of the computational box in the R direction, no absorbing mask is applied in this direction. The joint momentum distributions for electron and nuclei are obtained by using the mask method, see [42, 43]. The electron momentum distributions are calculated by integrating the joint momentum distributions over the proton momenta. The ground-state wave function of the model H_2^+ molecular ion is obtained by imaginary time propagation, providing the ground-state energy $E_0 = -0.78$ a.u., which is consistent with [39, 40].

Similar to [22], our first approach involves instantaneous excitation of the H_2^+ molecule. More specifically, the time propagation of the TDSE (1) starts with the wave function $\Psi_0(x, R) = \chi_0(R) \phi_1(x, R)$. Here $\phi_1(x, R)$ is the first excited electronic state for fixed R , and $\chi_0(R)$ is the nuclear ground-state wave function for the electronic ground state. Accordingly, the wave function $\phi_1(x, R)$ is a solution of the following time-independent Schrödinger equation

$$\left[-\frac{1}{2\mu} \frac{\partial^2}{\partial x^2} + V_{\text{pe}} + \frac{1}{\sqrt{R^2 + \alpha_1}} \right] \phi_n(x, R) = \varepsilon_n(R) \phi_n(x, R), \quad (6)$$

where the internuclear distance R is a parameter. In practice, we use imaginary time propagation for the time-dependent analogue of (6) to find $\phi_1(x, R)$. To enforce the correct symmetry, we symmetrize $\phi_0(x, R)$ [$1s\sigma_g$ state] and antisymmetrize the $\phi_1(x, R)$ [$2p\sigma_u$ state] at every step of the imaginary time propagation. The ground state energy $\varepsilon_0(R)$ is required to formulate the Schrödinger equation for the nuclear motion:

$$\left[-\frac{1}{m_p} \frac{\partial^2}{\partial R^2} + \varepsilon_0(R) \right] \chi_0(R) = \varepsilon_0 \chi_0(R). \quad (7)$$

Ionization of the H_2^+ molecule is performed by a strong and short probe pulse acting after a certain time delay t_c , see equation (3), that determines the internuclear distance at the ionization time. We first solve the TDSE (1) without the probe pulse [$A(t) = 0$] and calculate the average internuclear distance $\langle R \rangle$ as a function of time:

$$\langle R \rangle = \int dx \int dR R |\Psi(x, R, t_c)|^2. \quad (8)$$

Knowing this function allows us to assign an internuclear distance as a label to a PMD calculated for a given delay of the probe pulse. Then, the TDSE (1) is solved in the presence of the probe pulse, i.e. $A(t) = A_{\text{probe}}(t)$ to obtain the PMDs. To obtain the training data from fixed-nuclei calculations, we solve the 1D TDSE

$$i \frac{\partial}{\partial t} \Psi(x, t) = \left\{ -\frac{1}{2} \frac{\partial^2}{\partial x^2} - iA(t) \frac{\partial}{\partial x} - \frac{1}{\sqrt{(x - \frac{1}{2}R)^2 + \alpha_2}} - \frac{1}{\sqrt{(x + \frac{1}{2}R)^2 + \alpha_2}} \right\} \Psi(x, t) \quad (9)$$

for ionization from the first excited state at different values of the internuclear distance R .

As a second approach, we consider the well-known pump-probe scheme. More specifically, we solve the TDSE (1) with the vector potential given by $A(t) = A_{\text{pump}}(t) + A_{\text{probe}}(t)$,

$$A_{\text{pump}}(t) = \frac{F_{01}}{\omega_1} f_1(t) \sin(\omega_1 t), \quad (10)$$

where F_{01} and ω_1 are the field strength and the frequency of the pump pulse, respectively, and

$$f_1(t) = \begin{cases} \sin^2\left(\frac{\omega_1 t}{2n_{\text{p1}}}\right) & \text{for } 0 \leq t \leq \tau_{\text{pump}} \\ 0 & \text{otherwise.} \end{cases} \quad (11)$$

Here, n_{p1} and $\tau_{\text{pump}} = (2\pi/\omega_1) \cdot n_{\text{p1}}$ denote the number of cycles and the duration of the pump pulse, respectively. Since the envelope (11) reaches its maximum at $t = \pi n_{\text{p1}}/\omega_1$, the delay between the pump and probe pulses is given by $t_c - \pi n_{\text{p1}}/\omega_1$.

2.2. Architecture and training of neural networks

The PMDs obtained from the solution of the TDSE (with moving nuclei or fixed nuclei, see below) are used for training of neural networks. More specifically, we split the set of calculated PMDs into the training and test sets using the ratio 0.75 : 0.25. It should be noted that the resulting PMDs are functions of only one variable—the electron momentum component along the polarization direction. Therefore, application of a CNN is not the only option, and a fully connected neural network (FCNN) can also be applied to retrieve the bond length from PMDs. In the following, we shall use both the FCNN and CNN and compare the respective results with each other. We consider the PMDs in the range $-3.0 < k_x < 3.0$ a.u., and we normalize the distributions to their maximum values. As a result, every PMD is represented by a set of 765 values between 0 and 1. These sets are given to a neural network as input. In the present study, we employ two different FCNNs, see figures 1(a) and (b). Apart from an input layer, the first FCNN consists of only two fully connected layers with 765 and 1512 neurons, respectively, a dropout layer to prevent overfitting (serving as a regularization tool, see, e.g., [44]), and a last fully connected layer that predicts the output of the network, see figure 1(a). The number of neurons in this layer corresponds to the number of output quantities predicted by the network. The first two fully connected layers apply the rectified linear unit (ReLU) activation function: $\text{RELU}(x) = \max(0, x)$. The architecture of the second FCNN is shown in figure 1(b). While the first two fully connected layers of this network are similar to those of the first FCNN, it additionally includes a third layer with 1512 neurons, a fourth layer with 1512 neurons, and a fifth layer with 512 neurons. Consequently, this network contains a larger number of trainable parameters, about $7.1 \cdot 10^6$, compared with $1.75 \cdot 10^6$ trainable parameters of the network shown in figure 1(a). We use the MATLAB package [45] to implement the neural network. We split the training data into minibatches consisting of a certain number of distributions in order to use one minibatch for each iteration of the training process. The sizes of the minibatches used for training different FCNNs are specified in sections 3.1 and 3.2. The goal of training is to minimize the loss function (in our case the mean squared error), which characterizes the deviation between predictions of the neural network and the known labels of the training PMDs, i.e., internuclear distances. The training of the network starts using the learning rate $l_r = 5 \cdot 10^{-3}$, which is decreased by a factor of 10 after 20 training epochs. About 60 epochs are needed for convergence of the loss function. The training data are shuffled before each epoch, and therefore each PMD leads to an unbiased modification of the neural network. For training, we use the stochastic gradient descent (SGD) optimizer.

Similar to the networks used in studies [20–22], our present CNN consists of five pairs of convolutional layers and average pooling layers, dropout layer, and a fully connected layer calculating the output of the network, see figure 1(c). In contrast to CNNs [20–22], the pairs of convolutional and average pooling layers are preceded by a fully connected layer with $(2 \cdot 765) = 1530$ neurons. Both convolutional and average pooling layers work in only one dimension. Each convolutional layer consists of 32 filters, with each filter having a size of 2 points. Apart from the convolution of the input, the convolutional layers apply the ReLU activation function. The size of the pooling region in average pooling layers is also equal to 2 points. We find that for the problem at hand, good performance of the CNN with this architecture is achieved for a minibatch consisting of 30 images and the initial learning rate equal to 10^{-2} . The learning rate is again decreased by a factor of ten after 20 epochs. As with the FCNNs, the SGD optimizer is used for training the CNN.

3. Results and discussion

3.1. Retrieval of the time-dependent bond length for instantaneous excitation

For simplicity, we start the application of deep learning with the case of fixed laser intensity. We solve the TDSE (1) starting with instantaneous ionization and using $N = 310$ random delay times of the probe pulse and an intensity of $2.52 \times 10^{14} \text{ W cm}^{-2}$. The range of delay times corresponds to $3.61 < R < 12.00$ a.u. The resulting PMDs (see, e.g., figure 2(a)) were used to train a FCNN aimed at the retrieval of the internuclear distance at the ionization time. The loss function converges after about 40 training epochs, and the neural network recognizes the internuclear distance with a mean absolute error (MAE) of 0.07 a.u.

We then allow the intensity to vary in the range $[1.0, 4.0] \times 10^{14} \text{ W cm}^{-2}$, and we calculate $N = 2000$ PMDs for random internuclear distances and laser intensities. We train the FCNN shown in figure 1(a). The minibatches consist of 100 distributions. The neural network predicts the internuclear distance and the laser intensity with MAEs equal to 0.08 a.u. and 0.05 W cm^{-2} , respectively




Figure 1. The architecture of the (a), (b) FCNNs and (c) CNN used for the retrieval of the bond length R . Both neural networks take 1D electron momentum distributions normalized to their maximum value as input; see text.




Figure 2. Electron momentum distributions for ionization of the 1D H_2^+ molecule calculated from the direct numerical solution of the TDSE for the internuclear distance $R = 9.26$ a.u. The molecule is ionized by a laser pulse with duration $n_p = 2$ cycles, peak intensity 2.5×10^{14} W cm $^{-2}$, and wavelength 800 nm. (a) Momentum distribution obtained from the moving-nuclei TDSE for instantaneous excitation of the molecule and ionization at $\langle R \rangle = 9.26$ a.u. (b) Distribution calculated within the pump-probe scheme. The wavelength of the pump pulse is 148 nm ($\omega = 0.368$ a.u.), the peak laser intensity is 8.4×10^{12} W cm $^{-2}$, and duration of this pulse is $n_p = 3$ optical cycles. (c) Distribution obtained from the solution of the fixed-nuclei TDSE (9) for ionization of the ground state. (d) Distribution calculated from equation (9) for ionization of the first excited state. The distributions are normalized to their maximum value.




Figure 3. Plots of predicted vs true internuclear distances [(a),(c)] and laser intensities [(b),(d)] illustrating the performance of neural networks for the case of instantaneous excitation. (a) and (b) correspond to an FCNN trained on a set of distributions calculated from the TDSE (1) with moving nuclei. (c) and (d) show the performance of an FCNN trained on a set of PMDs with fixed nuclei, i.e., obtained from the solution of equation (9), for ionization from the first excited state. The dashed curves in (a) and (c) show the time-dependent expectation value of the bond length.

(see figures 3(a) and (b)). For comparison, in addition to the FCNN, we also train a CNN for this purpose. The CNN predicts R and I with the MAEs equal to 0.094 a.u. and 0.068 W cm^{-2} , respectively. Therefore, the use of the CNN does not provide an advantage over a FCNN for the problem at hand.

As a next step, neural networks trained on distributions calculated for fixed internuclear distances are applied to PMDs obtained for moving nuclei. A set of $N = 2000$ fixed-nuclei PMDs for ionization from the first excited state is used to train an FCNN (see figure 1(b)) to retrieve R and I . An example of such a distribution is shown in figure 2(d). This distribution should be compared with the PMD for ionization from the ground state shown in figure 2(c). The neural network shows good performance on a test set of PMDs with fixed nuclei: the corresponding MAEs for R and I are equal to 0.10 a.u. and 0.04 W cm^{-2} , respectively. However, its performance on distributions obtained for moving nuclei is considerably worse, although it remains acceptable: $\text{MAE}(R) = 0.48 \text{ a.u.}$, $\text{MAE}(I) = 0.12 \text{ W cm}^{-2}$, see figures 3(c) and (d). A CNN trained on the distributions for fixed nuclei shows similar performance. These results are easy to understand, considering that fixed nuclei PMDs differ from momentum distributions calculated for moving nuclei, see figures 2(a) and (c).

3.2. Retrieval of the time-dependent bond length in the pump-probe scheme

We perform the calculations for the pump pulse frequency $\omega = 0.3068 \text{ a.u.}$, which corresponds to the resonance between the σ_g and σ_μ states in our model molecule, the field strength $F_{01} = 0.0155 \text{ a.u.}$ (intensity of $8.4 \times 10^{12} \text{ W cm}^{-2}$), and $n_{p1} = 3$. As in section 3.1, we first calculate the time-dependent average internuclear distance $\langle R \rangle$ as a function of time in the absence of the probe pulse. This expectation value is calculated after projecting out the electronic-vibrational ground state at the end of the pump pulse [39]. We next solve the TDSE (1) with the vector potential (10) and we calculate a set of $N = 2000$ momentum distributions corresponding to various values of the time delay τ , and therefore various values of the internuclear distance $\langle R \rangle$, lying in the range $[3.98, 11.96] \text{ a.u.}$ An example of a PMD is shown in figure 2(b). Every PMD is labelled with an internuclear distance R in accordance with the function $\langle R \rangle(t)$. This set of PMDs is used for training of the FCNN shown in figure 1(b) and a CNN aimed at the recognition of the internuclear distance and the laser intensity. We find that the FCNN retrieves the internuclear distance with an MAE of 0.13 a.u. and the laser intensity with

an MAE of $0.05 \times 10^{14} \text{ W cm}^{-2}$. The CNN shows similar performance: $\text{MAE}(R) = 0.15 \text{ a.u.}$ and $\text{MAE}(I) = 0.044 \times 10^{14} \text{ W cm}^{-2}$.

At the next stage of our study, we apply neural networks trained on the distributions calculated for fixed nuclei to PMDs obtained in the pump-probe scheme. The FCNN shows poor performance in recognizing the internuclear distance when applied to PMDs obtained in the pump-probe scheme. The corresponding MAE for the internuclear distance is equal to 1.13 a.u., and the laser intensity can be reconstructed with an MAE of $0.25 \times 10^{14} \text{ W cm}^{-2}$. The CNN trained on PMDs with fixed nuclei demonstrates similar performance. These results are substantially worse compared to the case of instantaneous excitation (see section 3.1). In order to simplify the task of the neural network, we fix the laser intensity to $2.52 \times 10^{14} \text{ W cm}^{-2}$. We calculate a set of corresponding fixed nuclei PMDs, and we train an FCNN using this set. However, this fixed-intensity neural network performs only marginally better than the previous ones. For PMDs calculated for moving nuclei, it predicts the internuclear distance with an MAE of 1.03 a.u.

These results could have been expected. Indeed, PMDs calculated from the TDSE (1) in the pump-probe scheme for given values of $\langle R \rangle$ at the ionization time differ from the PMDs obtained from equation (9) for the same internuclear distances. This applies to ionization of both the ground and excited states in the TDSE (9) for fixed nuclei (see figure 2(b) with figures 2(c) and (d)). Analysis of the populations in the case of the TDSE (1) shows that the population of the ground electronic-vibrational state at the end of the pump pulse is equal to 0.92. This means that the pump pulse creates a superposition of the ground state (with a larger weight) and the first excited state (with a much smaller weight). This suggests that a set of PMDs defined as

$$\text{PMD}_i = w_i \text{PMD}_{\phi_{0,i}}(R_{0,i}) + (1 - w_i) \text{PMD}_{\phi_{1,i}}(R_{1,i}) \quad (12)$$

can be used for training of a fixed-nuclei neural network aimed at the retrieval of the time-varying bond length. Here i is the number of a distribution in the set, $\text{PMD}_{\phi_{0,i}}(R_{0,i})$ and $\text{PMD}_{\phi_{1,i}}(R_{1,i})$ are electron momentum distributions calculated from equation (9) for ionization of the ground and first excited states, respectively, and w_i is the weight of $\text{PMD}_{\phi_{0,i}}$. Note that the interatomic distances $R_{0,i}$ and $R_{1,i}$, for which the distributions $\text{PMD}_{0,i}$ and $\text{PMD}_{1,i}$ are calculated, do not necessarily have to be the same. Therefore, a set of PMDs (12) is determined by the choice of $R_{0,i}$, $R_{1,i}$, and w_i . The approximation of PMDs calculated from the non-Born–Oppenheimer TDSE (1) by the incoherent sum (12) is valid if the overlap between the nuclear wave packet of the excited state and the nuclear wave packet of the ground state is small. This condition is fulfilled provided that the time delay between the pump pulse and the probe pulse is large enough, which is a mild requirement for typical parameters.

We first fix R_0 to the equilibrium distance equal to 2.59 a.u. (for $\alpha_2 = 1.0$), and we calculate a set of $\text{PMD}_{\phi_1}(R_{1,i})$ for $R_{1,i}$ distributed randomly in the range $4.1 \leq R_{1,i} \leq 11.5 \text{ a.u.}$ The idea behind this choice is that only the wave packet in the excited electronic state undergoes dissociation. For simplicity, we again fix the laser intensity to $2.52 \cdot 10^{14} \text{ W cm}^{-2}$. The choice of the weights w_i is of crucial importance for the training procedure, and it turns out to be a challenging task. This is easy to understand if we compare the maximum of the $\text{PMD}_{\phi_0}(2.59 \text{ a.u.})$ equal to $5.93 \cdot 10^{-10}$ to the maxima of $\text{PMD}_{\phi_1}(R_1)$ for different values of R_1 . These latter maxima only weakly depend on R_1 . For example, $\max[\text{PMD}_{\phi_1}(R_1)]$ for $R_1 = 4.0 \text{ a.u.}$, $R_1 = 6.0 \text{ a.u.}$, and $R_1 = 8.0 \text{ a.u.}$ are equal to $2.96 \cdot 10^{-6}$, $2.89 \cdot 10^{-6}$, and $2.29 \cdot 10^{-6}$, respectively. The choice of the range for the weights w_i is based on the requirement that both terms in the sum (12) have comparable values. Specifically, we can assess the comparability, using the ratio $w \max[\text{PMD}_{\phi_0}(R_0)] / \{(1 - w) \max[\text{PMD}_{\phi_1}(R_1)]\}$. The range can be fine-tuned by adjusting, generating different sets of PMDs, training the corresponding neural networks, and evaluating their performance. Through this process, we have determined that optimal results are achieved for $w_i \in [0.99, 0.99999]$, i.e., for $(1 - w_i) \in [10^{-5}, 0.01]$. This suggests applying the neural network to a set of PMDs calculated for a lower intensity of the pump pulse than the intensity of $8.4 \times 10^{12} \text{ W cm}^{-2}$ used above. The pump-pulse intensity should be such that the ground-state population after the end of the pump pulse falls within the range of the weights used for training of the neural network. Therefore, we calculate another set of PMDs for the pump pulse intensity of $3.29 \times 10^{10} \text{ W cm}^{-2}$ corresponding to the field strength $F_{01} = 0.00097 \text{ a.u.}$ For this intensity, the ground-state population at the end of the pump pulse is 0.99948.

As is commonly acknowledged, neural network training is inherently stochastic, see, e.g., [36]. Different neural networks trained on the same training data set show differences in predicting internuclear distance. In the present case, this effect is particularly pronounced. For this reason, we train 10 different FCNNs on the same data set (12), which consists of 5000 distributions, and apply them to the test set of PMDs discussed above. The architecture of these neural networks is shown in figure 1(b), and




Figure 4. Predictions of neural networks for the internuclear distance during dissociation (blue points) at different time delays compared to the time-dependent expectation value of the internuclear distance (red dashed curves) obtained from the solution of the TDSE (1) for the pump-probe scheme. (a) Results from the FCNN trained on momentum distributions calculated from equation (12) for fixed $R_0 = 2.59$ a.u. The pump pulse intensity is $3.29 \times 10^{10} \text{ W cm}^{-2}$. (b) Results from the FCNN trained on the set of distributions (12) with $R_{0,i}$ randomly distributed in the range $[1.0, 3.0]$ a.u. The PMDs of the test set are again calculated for the intensity of the pump pulse equal to $3.29 \times 10^{10} \text{ W cm}^{-2}$.

the minibatch size is set to 60 images. We find that these neural networks retrieve the time-dependent internuclear distance with an MAE of 0.48 a.u. (averaged over neural networks). The performance of one selected FCNN is illustrated by figure 4(a). Simultaneously, the FCNNs show slightly better performance on PMDs calculated for higher pump-probe intensity. For example, for the intensity of $8.4 \times 10^{12} \text{ W cm}^{-2}$, the average MAE is equal to 0.40 a.u. This counterintuitive result along with the relatively low accuracy of the retrieved R suggests that the approach should be modified.

In our next approach, we vary not only R_1 but also R_0 in equation (12). More specifically, while the values $R_{1,i}$ vary again in the range $[4.1, 11.5]$ a.u., $R_{0,i}$ are randomly chosen from the range $[1.0, 3.0]$ a.u. It should be noted that $\max[\text{PMD}_{\phi_0}(R_0)]$ changes significantly with increasing R_0 . Indeed, $\max\text{PMD}_{\phi_0}(1.02) = 1.14 \cdot 10^{-10}$, $\max\text{PMD}_{\phi_0}(2.02) = 2.10 \cdot 10^{-10}$, and $\max\text{PMD}_{\phi_0}(3.02) = 1.74 \cdot 10^{-9}$. This makes the problem of identifying a suitable interval for w_i more difficult as compared to the previous case. Nevertheless, we find that the weights $w_i \in [0.999, 0.999999]$ provide the best results. For the pump-probe intensity of $3.29 \times 10^{10} \text{ W cm}^{-2}$, a set of $N = 10$ FCNNs with the architecture shown in figure 1(b) trained on 10000 PMDs (12) with random weights in this interval predicts the time-dependent internuclear distance $R_{1,i}$ with an averaged MAE of 0.19 a.u. Figure 4(b) shows the performance of one FCNN selected from this set.

We also apply the same method for the retrieval of the internuclear distances $R_{0,i}$ and weights w_i from pump-probe PMDs. In this case, the spread of predictions from different neural networks trained on the same data set is relatively small. The average internuclear distance $\langle R_{0,i} \rangle$ for the ground state delivered by a FCNN for the set of PMDs calculated for the pump pulse intensity of $3.29 \times 10^{10} \text{ W cm}^{-2}$ is equal to 2.68 a.u. This value is close to the equilibrium distance of 2.59 a.u. in the Born–Oppenheimer approximation. Simultaneously, the neural network predicts the average weight $\langle w_i \rangle$ of 0.99955. This value is close to the ground-state population 0.99948 at the end of the pump pulse. The obtained results show that the neural network trained on PMDs for frozen nuclei may be used for the retrieval of the bound-state populations and the internuclear distance in both dissociating and non-dissociating states. More research is needed to analyze the potential of this scheme in view of the fact that the ionization probability is strongly R -dependent even within the narrow range of the ground state nuclear distribution.

4. Conclusions

In conclusion, we have investigated the retrieval of the time-dependent internuclear distance of a dissociating 1D molecule using a deep neural network that takes PMDs produced by strong-field ionization as input. To this end, we either assumed an instantaneous excitation of the molecule to the first excited electronic state with subsequent dissociation or applied a pump-probe scheme. In the latter approach

the nuclear motion is initiated by a pump pulse that excites the molecule. The dissociating molecule is ionized by the probe pulse coming after a certain time delay, which determines the internuclear distance at the time of ionization. In both cases the corresponding PMDs were calculated from direct numerical solution of the TDSE with the non-Born–Oppenheimer treatment of the coupled electronic-nuclear motion. We have shown that a deep neural network (FCNN or CNN) trained on a few thousand PMDs retrieves the internuclear distance with an absolute error of about 0.1 a.u. This applies to both FCNNs and CNNs.

We have also applied a neural network trained on momentum distributions obtained at fixed internuclear distances. For instantaneous excitation of the molecule, the neural network retrieves the time-dependent internuclear distance with an MAE of 0.4 a.u. For application in the pump-probe scheme, a network has been trained on incoherent sums of electron momentum distributions calculated for ionization from the electronic ground state and from the first excited state with frozen nuclei taken with randomly distributed weights. We have found that this neural network can reconstruct the time-dependent internuclear distance with an absolute error of 0.2 a.u. This applies to both simulation schemes and both types of neural networks (FCNN and CNN) used in our study. Therefore, our results support the applicability of deep learning to the retrieval of the dissociation dynamics of molecules from electron momentum distributions.

In general, the accuracy of retrieving R using neural networks trained on PMDs for fixed nuclei achieved in the present study (0.2–0.4 a.u.) is slightly lower than the accuracy of 0.2–0.3 a.u. reported in [22]. We attribute this to two main factors. First, our current study inherently includes the averaging of PMDs over the range of internuclear distances covered by the nuclear wave packet. Second, we account for the nuclear motion during the probe pulse. Both these factors are absent in the fixed-nuclei PMDs at single internuclear distances used for training the neural networks. Consequently, we expect retrieval errors to be larger compared to the previous study, where these effects were included only partially. However, a thorough analysis of reconstruction errors deserves further studies as [22] analyzes the 2D H_2^+ molecule, while the current study concentrates on a 1D case.

The fact that neural networks trained on PMDs calculated for fixed nuclei can retrieve the internuclear distance based on distributions obtained by a complete quantum mechanical description of the molecule opens the way for application of the method to experimental data. Indeed, calculation of even a large set of PMDs from the 3D TDSE for a molecule within a frozen nuclei approximation is a feasible task. Applying the transfer learning technique to the neural network trained on this set of distributions will make the corresponding neural network applicable to focal-volume-averaged PMDs. Such a modified neural network is directly applicable to experimental PMDs. We note that the application of transfer learning does not require calculation of thousands of focal-volume-averaged momentum distributions: only a few hundreds of them are usually enough, see [20, 21].

Moreover, our work indicates that a neural network trained on PMDs with frozen nuclei may be used to reconstruct the populations of the electronic states. Therefore, this work demonstrates the application of deep learning to the retrieval of electronic properties in addition to nuclear motion. Advances in this direction are useful for the development of innovative tools for static and time-dependent molecular imaging.

Data availability statement

The data cannot be made publicly available upon publication because they are not available in a format that is sufficiently accessible or reusable by other researchers. The data that support the findings of this study are available upon reasonable request from the authors.

ORCID iDs

N I Shvetsov-Shilovski  0000-0002-8825-9302

M Lein  0000-0003-1489-8715

References

- [1] Carleo G, Cirac I, Cranmer K, Daudet L, Schuld M, Tishby N, Vogt-Maranto L and Zdeborová L 2019 Machine learning and the physical sciences *Rev. Mod. Phys.* **91** 045002

- [2] Suresh R, Bishnoi H, Kuklin A V, Parikh A, Molokeev M, Harinarayanan R, Gharat S and Hiba P 2024 Revolutionizing physics: a comprehensive survey of machine learning applications *Front. Phys.* **12** 1322162
- [3] Becker W, Grasbon F, Kopold R, Milošević D B, Paulus G G and Walther H 2002 Above-threshold ionization: from classical features to quantum effects *Adv. At. Mol. Opt. Phys.* **48** 35
- [4] Milošević D B and Ehlötzky F 2003 Scattering and reaction processes in powerful laser fields *Adv. At. Mol. Opt. Phys.* **49** 373
- [5] Becker A and Faisal F H 2005 Intense-field many-body S-matrix theory *J. Phys. B: At. Mol. Opt. Phys.* **38** R1
- [6] Figueira de Morisson Faria C and Liu X 2011 Electron–electron correlation in strong laser fields *J. Mod. Opt.* **58** 1076
- [7] M Kitzler and S Gräfe (eds) 2016 *Ultrafast Dynamics Driven by Intense Light Pulses* (From Atoms to Solids, From Lasers to Intense x-rays) (Springer) (<https://doi.org/10.1007/978-3-319-20173-3>)
- [8] Lin C D, Le A-T, Jin C and Wei H 2018 *Attosecond and Strong-Field Physics (Principles and Applications)* (Cambridge University Press) (<https://doi.org/10.1017/9781108181839>)
- [9] Gherman A M M, Kováčik K, Cristea M V and Toşa V 2018 Artificial neural network trained to predict high-harmonic flux *Appl. Sci.* **8** 2106
- [10] Zahavy T, Dikopoltsev A, Moss D, Haham G I, Cohen O, Mannor S and Segev M 2018 Deep learning reconstruction of ultrashort pulses *Optica* **5** 666
- [11] Kleinert S, Tajalli A, Nagy T and Morgner U 2019 Rapid phase retrieval of ultrashort pulses from dispersion scan traces using deep neural networks *Opt. Lett.* **44** 979
- [12] Liu X, Zhang G, Li J, Shi G, Zhou M, Huang B, Tang Y, Song X and Yang W 2020 Deep Learning for Feynman's path integral in strong-field time-dependent dynamics *Phys. Rev. Lett.* **124** 113002
- [13] Yan T M, Popruzhenko S V, Vrakking M J J and Bauer D 2010 Low-energy structures in strong field ionization revealed by quantum orbits *Phys. Rev. Lett.* **105** 253002
- [14] Yan T M and Bauer D 2012 Sub-barrier Coulomb effects on the interference pattern in tunneling-ionization photoelectron spectra *Phys. Rev. A* **86** 053403
- [15] Szoldra T, Ciappina M F, Werby N, Bucksbaum P H, Lewenstein M, Zakrzewski J and Maxwell A S 2023 Femtosecond pulse parameter estimation from photoelectron momenta using machine learning *New J. Phys.* **25** 083039
- [16] Pablos-Marín J M, Serrano J and Hernández-García C 2023 Simulating macroscopic high-order harmonic generation driven by structured laser beams using artificial intelligence *Comput. Phys. Commun.* **291** 108823
- [17] Lytova M, Spanner M and Tamblyn I 2023 Deep learning and high harmonic generation *Can. J. Phys.* **101** 132
- [18] Liu X, Amini K, Sanchez A, Belsa B, Steinle T and Biegert J 2021 Machine learning for laser-induced electron diffraction imaging of molecular structures *Commun. Chem.* **4** 154
- [19] He L *et al* 2022 Filming movies of attosecond charge migration in single molecules with high harmonic spectroscopy *Nat. Commun.* **13** 4595
- [20] Shvetsov-Shilovski N I and Lein M 2022 Deep learning for retrieval of the internuclear distance in a molecule from interference patterns in photoelectron momentum distributions *Phys. Rev. A* **105** L021102
- [21] Shvetsov-Shilovski N I and Lein M 2023 Transfer learning, alternative approaches and visualization of a convolutional neural network for retrieval of the internuclear distance in a molecule from photoelectron momentum distributions *Phys. Rev. A* **107** 033106
- [22] Shvetsov-Shilovski N I and Lein M 2024 Convolutional neural network for retrieval of the time-dependent bond length in a molecule from photoelectron momentum distributions *J. Phys. B: At. Mol. Opt. Phys.* **57** 06LT01
- [23] Xu J, Blaga C I, Agostini P and DiMauro L F 2016 Time-resolved molecular imaging *J. Phys. B: At. Mol. Opt. Phys.* **49** 112001
- [24] Meckel M *et al* 2008 Laser-induced electron tunneling and diffraction *Science* **320** 1748
- [25] Blaga C I, Xu J, DiChiara A D, Sistrunk E, Zhang K, Agostini P, Miller T A, DiMauro L F and Lin C D 2012 Imaging ultrafast molecular dynamics with laser-induced electron diffraction *Nature* **483** 194
- [26] Pullen M G *et al* 2015 Imaging an aligned polyatomic molecule with laser-induced electron diffraction *Nat. Commun.* **6** 7262
- [27] Huismans Y *et al* 2011 Time-resolved holography with photoelectrons *Science* **331** 61
- [28] De Giovannini U, Küpper J and Trabattoni A 2023 New perspectives in time-resolved laser-induced electron diffraction *J. Phys. B: At. Mol. Opt. Phys.* **56** 054002
- [29] Walt S G, Ram N B, Atala M, Shvetsov-Shilovski N, von Conta A, Baykusheva D, Lein M and Wörner H J 2017 Dynamics of valence-shell electrons and nuclei probed by strong-field holography and rescattering *Nat. Commun.* **8** 15651
- [30] Keil T and Bauer D 2017 Coulomb-corrected strong-field quantum trajectories beyond dipole approximation *J. Phys. B: At. Mol. Opt. Phys.* **50** 194002
- [31] Maurer J, Willenberg B, Daněk J, Mayer B W, Phillips C R, Gallmann L, Kläiber M, Hatsagortsyan K Z, Keitel C H and Keller U 2018 Probing the ionization wave packet and recollision dynamics with an elliptically polarized strong laser field in the nondipole regime *Phys. Rev. A* **97** 013404
- [32] He M, Li Y, Zhou Y, Li M, Cao W and Lu P 2018 Direct visualization of valence electron motion using strong-field photoelectron holography *Phys. Rev. Lett.* **120** 133204
- [33] He M, Zhou Y, Tan J, Li Y, Li M and Lu P 2018 Imaging charge migration in the asymmetric molecule with the holographic interference in strong-field tunneling ionization *J. Phys. B: At. Mol. Opt. Phys.* **51** 245602
- [34] Tan J, Zhou Y, He M, Ke Q, Liang J, Li Y, Li M and Lu P 2019 Time-resolved tunneling ionization via strong-field photoelectron holography *Phys. Rev. A* **99** 033402
- [35] Brennecke S and Lein M 2019 Strong-field photoelectron holography beyond the electric dipole approximation: a semiclassical analysis *Phys. Rev. A* **100** 023413
- [36] Goodfellow I, Bengio Y and Courville A 2016 *Deep Learning* (MIT)
- [37] Hiskes J R 1961 Dissociation of molecular ions by electric and magnetic fields *Phys. Rev.* **122** 1207
- [38] Kulander K C, Mies F H and Schafer K J 1996 Model for studies of laser-induced nonlinear processes in molecules *Phys. Rev. A* **53** 2562
- [39] Picón A, Bahabad A, Kapteyn H C, Murnane M M and Becker A 2011 Two-center interferences in photoionization of a dissociating H_2^+ molecule *Phys. Rev. A* **83** 013414
- [40] Mosert V and Bauer D 2015 Dissociative ionization of H_2^+ : Few-cycle effect in the joint electron-ion energy spectrum *Phys. Rev. A* **92** 043414
- [41] Feit M D, Fleck J A and Steiger A 1982 Solution of the Schrödinger equation by a spectral method *J. Comput. Phys.* **47** 412

- [42] Lein M, Marangos J P and Knight P L 2002 Electron diffraction in above-threshold ionization of molecules *Phys. Rev. A* **66** 051404(R)
- [43] Tong X M, Hino K and Toshima N 2006 Phase-dependent atomic ionization in few-cycle intense laser pulse *Phys. Rev. A* **74** 031405(R)
- [44] Trask A W 2019 *Grokking Deep Learning* (Manning)
- [45] The MathWorks Inc MATLAB Version 9.14 (R2023a) Natick, Massachusetts (available at: [mathworks.com](https://www.mathworks.com))